

BOSLEY MD REGULAR STRENGTH FOR WOMEN- minoxidil 2% solution
Scientific Hair Research, LLC

BosleyMD Regular Strength for Women Minoxidil Topical Solution 2%

Minoxidil 2% w/v

Hair regrowth treatment

USE

Use to regrow hair on the top of the scalp (vertex only, see picture on side of box)

For external use only.

Flammable: Keep away from fire or flame.

Do not use if:

- Your amount of hair loss is different than shown on the side of this carton, because this product may not work for you
- You have no family history of hair loss
- Your hair loss is sudden and/or patchy
- Your hair loss is associated with childbirth
- You do not know the reason for your hair loss
- You are under 18 years of age. Do not use on babies or children
- Your scalp is red, inflamed, infected, irritated, or painful
- You use other medicines on the scalp

Ask a doctor before use if you have heart disease

When using this product

- Do not apply on other parts of the body
- Avoid contact with the eyes. In case of accidental contact, rinse eyes with large amounts of cool tap water.
- Some people have experienced changes in hair color and/or texture
- It takes time to re-grow hair. You may need to use this product for at least 4 months before you see results.
- The amount of hair regrowth is different for each person. This product will not work for all women.

Stop use and ask a doctor if

- Chest pain, rapid heartbeat, faintness, or dizziness occurs
- Sudden, unexplained weight gain occurs
- Your hands or feet swell
- Scalp irritation or redness occurs
- Unwanted facial hair growth occurs
- You do not see hair regrowth in 4 months

May be harmful if used when pregnant or breast-feeding.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center immediately.


- Apply one mL (6 sprays) 2 times a day directly onto the scalp in the hair loss area

- Using more or more often will not improve results
- Continued use is necessary to increase and keep your hair regrowth, or hair loss will begin again
- See hair loss pictures on the side of this carton
- Before use, read all information on carton and enclosed leaflet
- Keep the carton. It contains important information
- In clinical studies of mostly white women aged 18-45 years with moderate degrees of hair loss, the following responses to Minoxidil Topical Solution 2% was reported: 19% of women reported moderate hair regrowth after using 2% minoxidil topical solution for 8 months (19% had moderate regrowth; 40% had minimal hair regrowth). This compares with the 7% of women reporting moderate hair regrowth after using a placebo, the liquid without minoxidil in it, for 8 months (7% had moderate regrowth; 33% had minimal regrowth).
- Store at controlled room temperature 20 ° to 25 ° C (68 ° to 77 °F)

Alcohol, Propylene Glycol, Purified Water

Questions or comments? Call your health professional or BosleyMD at 1(800)BOSLEY1

Mia Jenner



BOSLEY MD REGULAR STRENGTH FOR WOMEN

minoxidil 2% solution

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69113-004
Route of Administration	TOPICAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
MINOXIDIL (UNII: 5965120SH1) (MINOXIDIL - UNII:5965120SH1)		MINOXIDIL	2 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69113-004-01	2 in 1 BOX	05/21/2020	
1		60 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA078176	05/21/2020	

Labeler - Scientific Hair Research, LLC (961897647)

Revised: 5/2020

Scientific Hair Research, LLC